

**Mineralogical study of the seafloor diagenesis of carbonate sediments
near Dry Tortugas, Gulf of Mexico,
using Rietveld method of crystal structure refinement**

by

Yoko Furukawa


Center for Marine Sciences
The University of Southern Mississippi
Stennis Space Center, MS 39529

Quarterly Technical Report

Submitted to

Naval Research Laboratory (NRL-SSC)
Code 7431
Stennis Space Center, MS 39529-5004

Grant No. N00014-95-1-G907

Approved: 
Denis Wiesenburg, Principal Investigator

Technical Report No. CMS-95-05

19960301 040

December 31, 1995

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED

DTIC QUALITY INSPECTED 1

1. Introduction

Sediment microstructure is largely determined by chemical diagenesis and physical processes such as bioturbation and storm reworking in the carbonate sediments of Fort Jefferson National Monument area, west of Key West, Florida. Consequently, understanding of the extent of chemical diagenesis and physical reworking is essential in order to understand the sediment structure as well as the geophysical and geoacoustic properties affected by the sediment structure. This on-going study investigates the mineralogy, porewater geochemistry, and sediment microstructure of a number of box and gravity cores from the study area in order to quantify the extent of chemical diagenesis. This quarterly technical report summarizes the results of mineralogical study, whereas preliminary results of geochemical and microfabric study are presented in the previous technical reports (Furukawa, 1995a, b, c)

This study, funded by the Naval Research Laboratory (NRL), is a part of the large research program managed by the Office of Naval Research (ONR) called Coastal Benthic Boundary Layer Special Research Program (CBBL-SRP). The objective of the CBBL-SRP is to characterize and model benthic boundary layer processes and the impact such processes cause on seafloor properties that affect shallow-water naval operation (Richardson, 1995). The present study investigates the chemical diagenesis in an attempt to understand the relationship between sediment structure and processes that form the structure, chemical diagenesis, sedimentation and physical reworking.

2. Methods

The author participated in the CBBL-SRP Key West Campaign that was conducted in the area of carbonate sediments near Dry Tortugas (Fort Jefferson National Monument) and Marquesas Keys, west of Key West, Florida in the Gulf of Mexico. Dry Tortugas area sediments were sampled extensively to study mineralogy. Results from several box cores (KW-PL-BC-141, 165, 194), a diver core (KW-PL-DC-178), and a gravity core (KW-PE-GC-147) are reported herein. Sediments near Marquesas Keys were collected also and results from a box core (KW-PL-BC-208) are reported herein. The same box, diver and gravity cores were subsampled also for pore water chemistry and microfabric. The chemistry and microfabric results are presented in previous technical reports (Furukawa, 1995a, b, c).

Mineralogy is studied using X-ray powder diffraction (XRD) and the subsequent Rietveld method of crystal structure refinement (Rietveld, 1969; Young, 1995; Young et al., 1994). For some samples (KW-PE-GC-147), clay, silt, and sand fractions were separated using settling and siphoning in order to study the size-fraction mineralogy. The Rietveld method of refinement is capable of quantifying the calcium carbonate phases with various Mg contents (Bish and Post 1993). The Rietveld method also provides the cell constants of high-Mg and low-Mg calcites from which Mg contents can be calculated using the existing correlation curve (Goldsmith et al., 1958; Reid et al., 1993).

3. Results

The XRD profiles of bulk samples from the box and diver cores are shown in Figures 1-5. The XRD profiles of the gravity core are shown in Figure 6 for the bulk samples, and in Figures 7-9 for sand, silt and clay fraction samples. The peaks indicate that the sediments are a mixture of aragonite, high-Mg calcite, low-Mg calcite, and non-carbonate fraction that is dominated by quartz. Note the relative intensities of high-Mg calcite and low-Mg calcite peaks. Whereas the relative intensities remain nearly constant through the box and gravity core sampling depths, they change significantly at depths in clay fraction of the gravity core samples.

The Rietveld method of crystal structure refinement quantified the change in relative amount of high-Mg calcite (HMC) and low-Mg calcite (LMC) in clay, silt, and sand fractions of samples from KW-PE-GC-147 (Figure 10). The HMC/LMC ratio increases at depths in the clay fraction.

4. Discussion

The increase in HMC/LMC calcite ratio in a gravity core below 120 cm (Figure 10) may be the result of cementation. Most cement in shallow water sediments is either high-Mg calcite or aragonite (Bathurst, 1975). Because the effect of cementation should be more notable within the matrix than around the surface of grains, due to the surface area difference, the comparison of matrix (i.e., clay) mineralogy vs. grain (i.e., silt and sand) mineralogy clarifies the reason for the observed shift. Because the observed shift in HMC/LMC ratio comes from cementation, the clay mineralogy shows the HMC/LMC increase at depth whereas the grain mineralogy remains constant. If the shift had been from the difference in depositional environment that results in more primary HMC grains (e.g., red algae and benthic foraminefera) at depth, both matrix mineralogy and grain mineralogy would have shown the shift. The lack of shift within the upper 120 cm of the sediments may be due to the physical reworking and/or lack of cementation. Cementation may become significant right below the mixing layer, or may not be detectable by mineralogy until the sediment is buried without mixing for a long period of time.

5. Further study

More quantitative mineralogy using the Rietveld method is planned. Further microfabric study using scanning electron microscope (SEM) is underway. The results will be combined with mineralogy and pore water chemistry and presented at Mississippi Academy of Sciences meeting in February 1996.

6. References

- Bathurst, R. G. C. (1975) *Carbonate sediments and their diagenesis, Second Enlarged Edition*. Developments in Sedimentology, Vol. 12. Elsevier.

- Bish, D. L. and Post, J. E. (1993) Quantitative mineralogical analysis using the Rietveld full-pattern fitting method. *Amer. Mineral.* **78**, 932-940.
- Furukawa, Y. (1995a) Oxidation of aqueous sulfide in porewater as the possible cause of carbonate dissolution during early diagenesis: A progress report. Center for Marine Sciences Report CMS-95-01. The University of Southern Mississippi, 6 pp.
- Furukawa, Y. (1995b) Evaluation of early diagenesis in modern, shallow-water carbonate sediments by mineralogy, fabric and porewater geochemistry. Center for Marine Sciences Report CMS-95-02. The University of Southern Mississippi, 10 pp.
- Furukawa, Y. (1995c) Geochemical, mineralogical and microstructural study of the seafloor diagenesis of carbonate sediments in Fort Jefferson National Monument area. Center for Marine Sciences Report CMS-95-04. The University of Southern Mississippi, 15 pp.
- Goldsmith, J. R. and Graf, D. L. (1958) Relation between lattice constants and composition of the Ca-Mg carbonates. *Amer. Mineral.* **43**, 84-101.
- Richardson, M. (1995) Introduction. *Key West Campaign Cruise Reports* (compiled M. D. Richardson and L. Cole), Chap. 1. Naval Research Laboratory.
- Reid, R. P., MacIntyre, I. G. and Post, J. E. (1993) Micritized skeletal grains in northern Belize Lagoon: A major source of Mg-calcite mud. *J. Sed. Petrol.* **62**, 145-156.
- Rietveld, H. M. (1969) A profile refinement method for nuclear and magnetic structures. *J. Appl. Cryst.* **2**, 65-71.
- Young, R. A. (1995) The Rietveld Method, IUCr Monograph on Crystallography No. 5. International Union of Crystallography, Oxford Science Publications, United Kingdom.
- Young, R. A., Sakthivel, A., Moss, T. S., and Pavia-Santos, C. O. (1994) User's guide to program DBWS9411 for Rietveld analysis of X-ray and neutron powder diffraction patterns with a 'PC' and various other computers. School of Physics, Georgia Institute of Technology, Atlanta, Georgia.




Figure 1. X-ray powder diffraction profiles of bulk sediment samples from KW-PL-BC-141. Sample depths are indicated on the right in terms of centimeters below seafloor. AR, aragonite; LMC, low-Mg calcite; HMC, high-Mg calcite; QZ, quartz.




Figure 2. X-ray powder diffraction profiles of bulk sediment samples from KW-PL-BC-165. Sample depths are indicated on the right in terms of centimeters below seafloor. AR, aragonite; LMC, low-Mg calcite; HMC, high-Mg calcite; QZ, quartz.




Figure 3. X-ray powder diffraction profiles of bulk sediment samples from KW-PL-DC-178. Sample depths are indicated on the right in terms of centimeters below seafloor. AR, aragonite; LMC, low-Mg calcite; HMC, high-Mg calcite; QZ, quartz.




Figure 4. X-ray powder diffraction profiles of bulk sediment samples from KW-PL-BC-194. Sample depths are indicated on the right in terms of centimeters below seafloor. AR, aragonite; LMC, low-Mg calcite; HMC, high-Mg calcite; QZ, quartz.




Figure 5. X-ray powder diffraction profiles of bulk sediment samples from KW-PL-BC-208. Sample depths are indicated on the right in terms of centimeters below seafloor. AR, aragonite; LMC, low-Mg calcite; HMC, high-Mg calcite; QZ, quartz.




Figure 6. X-ray powder diffraction profiles of bulk sediment samples from KW-PE-GC-147.




Figure 7. X-ray powder diffraction profiles of sand fraction samples from KW-PE-GC-147.




Figure 8. X-ray powder diffraction profiles of silt fraction samples from KW-PE-GC-147.




Figure 9. X-ray powder diffraction profiles of clay fraction samples from KW-PE-GC-147.




Figure 10. HMC/LMC ratio of size-fractioned samples from KW-PE-GC-147.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. Agency Use Only (Leave blank).		2. Report Date. December 1995		3. Report Type and Dates Covered. Technical Report 10/95-12/95
4. Title and Subtitle. Mineralogical study of the seafloor diagenesis of carbonate sediments near Dry Tortugas, Gulf of Mexico, using Rietveld method of crystal structure refinement			5. Funding Numbers. Program Element No. Project No. Task No. Accession No.	
6. Author(s) Yoko Furukawa				
7. Performing Organization Name(s) and Address(es). Center for Marine Sciences The University of Southern Mississippi Building 1103, Room 102 Stennis Space Center, MS 39529			8. Performing Organization Report Number CMS-95-05	
9. Sponsoring/Monitoring Agency Name(s) and Address(es). Department of The Navy Naval Research Laboratory (NRL-SSC) CODE 3250: BJO Stennis Space Center, MS 39529			10. Sponsoring/Monitoring Agency Report Number. CR/7431--96-0001	
11. Supplementary Notes. ONR Research Grant No. N00014-95-1-G907				
12a. Distribution/Availability Statement. Approved for public releases; distribution is unlimited.			12b. Distribution Code. .	
13. Abstract (Maximum 200 words). The carbonate sediments from Fort Jefferson National Monument area were studied for mineralogy using X-ray powder diffraction (XRD) in order to evaluate the effects of seafloor diagenesis on sediment structure. Cementation is suggested at the depths of 120 cm and below by the increase in high-Mg/low-Mg calcite ratio within fine-grained matrix. The relative amounts of calcium carbonate phases were determined by the Rietveld crystal structure refinement. Seafloor diagenesis may be significant at the sediment depths which are not affected by bioturbation and storm mixing.				
14. Subject Terms. Seafloor diagenesis, Rietveld crystal structure refinement, carbonate mineralogy, Gulf of Mexico			15. Number of Pages. 15	
			16. Price Code.	
17. Security Classification of Report Unclassified	18. Security Classification of This Page. Unclassified	19. Security Classification of Abstract. Unclassified	20. Limitation of Abstract. SAR	